

Number Theory

Arithmetic Algebraic Geometry

11Gxx

- [1] Amod Agashe, Kenneth Ribet, and William A. Stein. The Manin constant. *Pure Appl. Math. Q.*, 2(2):617–636, 2006.
- [2] Amod Agashe and William Stein. Visibility of Shafarevich-Tate groups of abelian varieties. *J. Number Theory*, 97(1):171–185, 2002.
- [3] Amod Agashe and William Stein. Visible evidence for the Birch and Swinnerton-Dyer conjecture for modular abelian varieties of analytic rank zero. *Math. Comp.*, 74(249):455–484 (electronic), 2005.
- [4] Scott Ahlgren and Matthew Papanikolas. Higher Weierstrass points on $X_0(p)$. *Trans. Amer. Math. Soc.*, 355(4):1521–1535 (electronic), 2003.
- [5] Avner Ash, Darrin Doud, and David Pollack. Galois representations with conjectural connections to arithmetic cohomology. *Duke Math. J.*, 112(3):521–579, 2002.
- [6] Matthew H. Baker, Enrique González-Jiménez, Josep González, and Bjorn Poonen. Finiteness results for modular curves of genus at least 2. *Amer. J. Math.*, 127(6):1325–1387, 2005.
- [7] Arthur Baragar and Ronald van Luijk. $K3$ surfaces with Picard number three and canonical vector heights. *Math. Comp.*, 76(259):1493–1498 (electronic), 2007.
- [8] Mark Bauer, Edlyn Teske, and Annegret Weng. Point counting on Picard curves in large characteristic. *Math. Comp.*, 74(252):1983–2005 (electronic), 2005.
- [9] Tobias Berger and Krzysztof Klosin. A deformation problem for Galois representations over imaginary quadratic fields. *J. Inst. Math. Jussieu*, 8(4):669–692, 2009.
- [10] Amnon Besser and Rob De Jeu. $\text{li}(p)$ -service? an algorithm for computing p -adic polyalgorithms. *Math. Comp.*, 77(262):1105–1134, 2008.

- [11] Peter Birkner. *Efficient Arithmetic on Low-genus Curves*. Ph D thesis, Technische Universiteit Eindhoven, 2009.
- [12] Nigel Boston and Rafe Jones. Arboreal Galois representations. *Geom. Dedicata*, 124:27–35, 2007.
- [13] M. J. Bright, N. Bruin, E. V. Flynn, and A. Logan. The Brauer-Manin obstruction and Sh[2]. *LMS J. Comput. Math.*, 10:354–377 (electronic), 2007.
- [14] David Brown. The Chabauty-Coleman bound at a prime of bad reduction. [arXiv:0803.0973v2](https://arxiv.org/abs/0803.0973v2) [math.NT], 10 pages, 2008.
- [15] Ezra Brown and Bruce T. Myers. Elliptic curves from Mordell to Diophantus and back. *Amer. Math. Monthly*, 109(7):639–649, 2002.
- [16] N. Bruin and E. V. Flynn. n -covers of hyperelliptic curves. *Math. Proc. Cambridge Philos. Soc.*, 134(3):397–405, 2003.
- [17] Nils Bruin. Visualising *Sha*[2] in abelian surfaces. *Math. Comp.*, 73(247):1459–1476 (electronic), 2004.
- [18] Nils Bruin. The arithmetic of Prym varieties in genus 3. *Compos. Math.*, 144(2):317–338, 2008.
- [19] Nils Bruin and Noam D. Elkies. Trinomials $ax^7 + bx + c$ and $ax^8 + bx + c$ with Galois groups of order 168 and $8 \cdot 168$. In *Algorithmic Number Theory (Sydney, 2002)*, volume 2369 of *Lecture Notes in Comput. Sci.*, pages 172–188. Springer, Berlin, 2002.
- [20] Nils Bruin and E. Victor Flynn. Towers of 2-covers of hyperelliptic curves. *Trans. Amer. Math. Soc.*, 357(11):4329–4347 (electronic), 2005.
- [21] Nils Bruin and Michael Stoll. Deciding existence of rational points on curves: an experiment. *Experiment. Math.*, 17(2):181–189, 2008.
- [22] Nils Bruin and Michael Stoll. Two-cover descent on hyperelliptic curves. [arXiv:0803.2052v1](https://arxiv.org/abs/0803.2052v1) [math.NT], 19 pages, 2008.
- [23] Yann Bugeaud, Maurice Mignotte, Samir Siksek, Michael Stoll, and Szabolcs Tengely. Integral points on hyperelliptic curves. *Algebra Number Theory*, 2(8):859–885, 2008.

- [24] Kevin Buzzard and L. J. P. Kilford. The 2-adic eigencurve at the boundary of weight space. *Compos. Math.*, 141(3):605–619, 2005.
- [25] Robert Carls. Theta null points of 2-adic canonical lifts. [arXiv:math.NT/0509092](#), 18 pages, 2005.
- [26] Robert Carls and David Lubicz. A p -adic quasi-quadratic time point counting algorithm. *Int. Math. Res. Not. IMRN*, (4):698–735, 2009.
- [27] Antoine Chambert-Loir. Compter (rapidement) le nombre de solutions d'équations dans les corps finis. [arXiv:math.NT/0611584](#), 46 pages, 2006.
- [28] Denis Charles and Kristin Lauter. Computing modular polynomials. *LMS J. Comput. Math.*, 8:195–204 (electronic), 2005.
- [29] Denis Xavier Charles. Complex multiplication tests for elliptic curves. [arXiv:math.NT/0409501 v1](#), 13 pages, 2004.
- [30] Imin Chen. On the equation $s^2 + y^{2p} = \alpha^3$. *Math. Comp.*, 77(262):1223–1227, 2008.
- [31] Imin Chen and Chris Cummins. Elliptic curves with nonsplit mod 11 representations. *Math. Comp.*, 73(246):869–880 (electronic), 2004.
- [32] Robert F. Coleman and William A. Stein. Approximation of eigenforms of infinite slope by eigenforms of finite slope. In *Geometric Aspects of Dwork Theory. Vol. I, II*, pages 437–449. Walter de Gruyter GmbH & Co. KG, Berlin, 2004.
- [33] B. Conrad, K. Conrad, and H. Helfgott. Root numbers and ranks in positive characteristic. *Adv. Math.*, 198(2):684–731, 2005.
- [34] Brian Conrad, Bas Edixhoven, and William Stein. $J_1(p)$ has connected fibers. *Doc. Math.*, 8:331–408 (electronic), 2003.
- [35] Caterina Consani and Jasper Scholten. Arithmetic on a quintic threefold. *Internat. J. Math.*, 12(8):943–972, 2001.
- [36] Patrick Kenneth Corn. *Del Pezzo Surfaces and the Brauer-Manin Obstruction*. PhD Thesis, University of California, Berkley, 1998.

- [37] J. E. Cremona. *Algorithms for Modular Elliptic Curves*. Cambridge University Press, Cambridge, second edition, 1997.
- [38] J. E. Cremona, T. A. Fisher, C. O’Neil, D. Simon, and M. Stoll. Explicit n -descent on elliptic curves. I. Algebra. *J. Reine Angew. Math.*, 615:121–155, 2008.
- [39] J. E. Cremona and M. P. Lingham. Finding all elliptic curves with good reduction outside a given set of primes. *Experiment. Math.*, 16(3):303–312, 2007.
- [40] J. E. Cremona, M. Prickett, and Samir Siksek. Height difference bounds for elliptic curves over number fields. *J. Number Theory*, 116(1):42–68, 2006.
- [41] John Cremona. The elliptic curve database for conductors to 130000. In *Algorithmic number theory*, volume 4076 of *Lecture Notes in Comput. Sci.*, pages 11–29. Springer, Berlin, 2006.
- [42] John Cremona, Tom Fisher, Cathy O’Neil, Denis Simon, and Michael Stoll. Explicit n -descent on elliptic curves II: Geometry. [arXiv:math.NT/0611606](https://arxiv.org/abs/math.NT/0611606), 24 pages, 2006.
- [43] John E. Cremona. A solution for note 84.35. *The Mathematical Gazette*, 86(505):66–68, 2002.
- [44] John Cullinan. A computational approach to the 2-torsion structure of abelian threefolds. *Math. Comp.*, 78(267):1825–1836, 2009.
- [45] C. J. Cummins and S. Pauli. Congruence subgroups of $\mathrm{PSL}(2, Z)$ of genus less than or equal to 24. *Experiment. Math.*, 12(2):243–255, 2003.
- [46] Samit Dasgupta. Computations of elliptic units for real quadratic fields. *Canad. J. Math.*, 59(3):553–574, 2007.
- [47] Chantal David and Tom Weston. Local torsion on elliptic curves and the deformation theory of Galois representations. *Math. Res. Lett.*, 15(3):599–611, 2008.
- [48] Lassina Dembélé. A non-solvable galois extension of q ramified at 2 only. *C. R., Math., Acad. Sci. Paris*, 347(3-4), 2009.

- [49] Jan Denef and Frederik Vercauteren. An extension of Kedlaya’s algorithm to Artin-Schreier curves in characteristic 2. In *Algorithmic Number Theory (Sydney, 2002)*, volume 2369 of *Lecture Notes in Comput. Sci.*, pages 308–323. Springer, Berlin, 2002.
- [50] Xavier Charles Denis. Complex multiplication tests for elliptic curves. *Preprint*, 13 pages, 2004.
- [51] Claus Diem and Emmanuel Thomé. Index calculus in class groups of non-hyperelliptic curves of genus three. *J. Cryptology*, 21(4):593–611, 2008.
- [52] T. Dokchitser and V. Dokchitser. Computations in non-commutative Iwasawa theory. *Proc. Lond. Math. Soc. (3)*, 94(1):211–272, 2007.
- [53] Tim Dokchitser and Vladimir Dokchitser. Root numbers of elliptic curves in residue characteristic 2. *Bull. Lond. Math. Soc.*, 40(3):516–524, 2008.
- [54] Darrin Doud. A procedure to calculate torsion of elliptic curves over \mathbf{Q} . *Manuscripta Math.*, 95(4):463–469, 1998.
- [55] Andrej Dujella. On Mordell-Weil groups of elliptic curves induced by Diophantine triples. *Glas. Mat. Ser. III*, 42(62)(1):3–18, 2007.
- [56] S. Duquesne. Rational points on hyperelliptic curves and an explicit Weierstrass preparation theorem. *Manuscripta Math.*, 108(2):191–204, 2002.
- [57] Sylvain Duquesne. Points rationnels et méthode de Chabauty elliptique. *J. Théor. Nombres Bordeaux*, 15(1):99–113, 2003.
- [58] Sylvain Duquesne. Elliptic curves associated with simplest quartic fields. *J. Théor. Nombres Bordeaux*, 19(1):81–100, 2007.
- [59] Sylvain Duquesne. Montgomery ladder for all genus 2 curves in characteristic 2. In *Arithmetic of Finite Fields*, volume 5130 of *Lecture Notes in Computer Science*, pages 174–188. Springer, 2008.
- [60] Bas Edixhoven. On the computation of the coefficients of a modular form. In *Algorithmic number theory*, volume 4076 of *Lecture Notes in Comput. Sci.*, pages 30–39. Springer, Berlin, 2006.

- [61] Kirsten Eisentraeger, Dimitar Jetchev, and Kristin Lauter. On the computation of the Cassels pairing for certain Kolyvagin classes in the Shafarevich-Tate group. *5209:113–125*, 2008.
- [62] Kirsten Eisenträger and Kristin Lauter. A CRT algorithm for constructing genus 2 curves over finite fields. [arXiv:math.NT/0405305v2](https://arxiv.org/abs/math.NT/0405305v2), 16 pages, 2007.
- [63] Noam D. Elkies. Three lectures on elliptic surfaces and curves of high rank. [arXiv:0709.2908v1](https://arxiv.org/abs/0709.2908v1), 14 pages, 2007.
- [64] Noam D. Elkies. Shimura curve computations via K3 surfaces of Neron-Severi rank at least 19. In *Algorithmic Number Theory*, volume 5011 of *Lecture Notes in Computer Science*, pages 196–211. Springer, 2008.
- [65] Noam D. Elkies and Mark Watkins. Elliptic curves of large rank and small conductor. In *Algorithmic Number Theory*, volume 3076 of *Lecture Notes in Comput. Sci.*, pages 42–56. Springer, Berlin, 2004.
- [66] Arsen Elkin. Hyperelliptic Jacobians with real multiplication. *J. Number Theory*, 117(1):53–86, 2006.
- [67] Andreas-Stephan Elsenhans and Jörg Jahnel. K3 surfaces of Picard rank one and degree two. In *Algorithmic Number Theory*, volume 5011 of *Lecture Notes in Computer Science*, pages 212–225. Springer, 2008.
- [68] G. Everest and T. Ward. The canonical height of an algebraic point on an elliptic curve. *New York J. Math.*, 6:331–342 (electronic), 2000.
- [69] Graham Everest, Patrick Ingram, Valéry Mahé, and Shaun Stevens. The uniform primality conjecture for elliptic curves. *Acta Arith.*, 134(2):157–181, 2008.
- [70] Graham Everest, Patrick Ingram, and Shaun Stevens. Primitive divisors on twists of Fermat’s cubic. *LMS J. Comput. Math.*, 12:54–81, 2009.
- [71] Graham Everest and Valery Mahe. A generalization of Siegel’s theorem and Hall’s conjecture. *Exp. Math.*, 18(1):1–10, 2009.

- [72] Xander Faber and Benjamin Hutz. On the number of rational iterated pre-images of the origin under quadratic dynamical systems. [arXiv:0810.1715](https://arxiv.org/abs/0810.1715), 18 pages, 2008.
- [73] Reza Rezaeian Farashahi and Ruud Pellikaan. The quadratic extension extractor for (hyper)elliptic curves in odd characteristic. In *Arithmetic of finite fields*, volume 4547 of *Lecture Notes in Comput. Sci.*, pages 219–236. Springer, Berlin, 2007.
- [74] Julio Fernández, Josep González, and Joan-C. Lario. Plane quartic twists of $X(5, 3)$. *Canad. Math. Bull.*, 50(2):196–205, 2007.
- [75] Luís R. A. Finotti. Degrees of the elliptic Teichmüller lift. *J. Number Theory*, 95(2):123–141, 2002.
- [76] Luís R. A. Finotti. Minimal degree liftings of hyperelliptic curves. *J. Math. Sci. Univ. Tokyo*, 11(1):1–47, 2004.
- [77] Luís R. A. Finotti. Minimal degree liftings in characteristic 2. *J. Pure Appl. Algebra*, 207(3):631–673, 2006.
- [78] Tom Fisher. The Hessian of a genus one curve. [arXiv:math.NT/0610403](https://arxiv.org/abs/math.NT/0610403), 28 pages, 2006.
- [79] Tom Fisher. Testing equivalence of ternary cubics. In *Algorithmic Number Theory (Berlin, 2006)*, volume 4076 of *Lecture Notes in Comput. Sci.*, pages 333–345. Springer, Berlin, 2006.
- [80] Tom Fisher. A new approach to minimising binary quartics and ternary cubics. *Math. Res. Lett.*, 14(4):597–613, 2007.
- [81] Tom Fisher. The invariants of a genus one curve. *Proc. Lond. Math. Soc. (3)*, 97(3):753–782, 2008.
- [82] E. V. Flynn. The Hasse principle and the Brauer-Manin obstruction for curves. *Manuscripta Math.*, 115(4):437–466, 2004.
- [83] E. V. Flynn and C. Grattoni. Descent via isogeny on elliptic curves with large rational torsion subgroups. *J. Symbolic Comput.*, 43(4):293–303, 2008.

- [84] E. V. Flynn and J. Wunderle. Cycles of covers. *Monatsh. Math.*, Online first:16, 2008.
- [85] David Freeman, Peter Stevenhagen, and Marco Streng. Abelian varieties with prescribed embedding degree. In *Algorithmic Number Theory*, volume 5011 of *Lecture Notes in Computer Science*, pages 60–73. Springer, 2008.
- [86] S. D. Galbraith, J. F. McKee, and P. C. Valença. Ordinary abelian varieties having small embedding degree. *Finite Fields Appl.*, 13(4):800–814, 2007.
- [87] Steven D. Galbraith. Weil descent of Jacobians. *Discrete Appl. Math.*, 128(1):165–180, 2003.
- [88] P. Gaudry and É. Schost. Modular equations for hyperelliptic curves. *Math. Comp.*, 74(249):429–454 (electronic), 2005.
- [89] Pierrick Gaudry. Index calculus for abelian varieties and the elliptic curve discrete logarithm problem. *Preprint*, 13 pages, 2004.
- [90] Pierrick Gaudry and Nicolas Gürel. An extension of Kedlaya’s point-counting algorithm to superelliptic curves. In *Advances in Cryptology—Asiacrypt 2001 (Gold Coast)*, volume 2248 of *Lecture Notes in Comput. Sci.*, pages 480–494. Springer, Berlin, 2001.
- [91] Pierrick Gaudry and Robert Harley. Counting points on hyperelliptic curves over finite fields. In *Algorithmic Number Theory (Leiden, 2000)*, volume 1838 of *Lecture Notes in Comput. Sci.*, pages 313–332. Springer, Berlin, 2000.
- [92] Eknath Ghate, Enrique González-Jiménez, and Jordi Quer. On the Brauer class of modular endomorphism algebras. *Int. Math. Res. Not.*, (12):701–723, 2005.
- [93] Jean Gillibert. Invariants de classes: exemples de non-annulation en dimension supérieure. *Math. Ann.*, 338(2):475–495, 2007.
- [94] Edray Goins. Explicit descent via 4-isogeny on an elliptic curve. [arXiv:math.NT/0411215](https://arxiv.org/abs/math.NT/0411215) v1, 20 pages, 2004.

- [95] Josep González and Jordi Guàrdia. Genus two curves with quaternionic multiplication and modular Jacobian. *Math. Comp.*, 78(265):575–589, 2009.
- [96] Josep González, Jordi Guàrdia, and Victor Rotger. Abelian surfaces of GL_2 -type as Jacobians of curves. *Acta Arith.*, 116(3):263–287, 2005.
- [97] Josep González and Victor Rotger. Non-elliptic Shimura curves of genus one. *J. Math. Soc. Japan*, 58(4):927–948, 2006.
- [98] Enrique González-Jiménez, Josep González, and Jordi Guàrdia. Computations on modular Jacobian surfaces. In *Algorithmic Number Theory (Sydney, 2002)*, volume 2369 of *Lecture Notes in Comput. Sci.*, pages 189–197. Springer, Berlin, 2002.
- [99] Eyal Z. Goren and Kristin E. Lauter. The distance between superspecial abelian varieties with real multiplication. *J. Number Theory*, 129(6):1562–1578, 2009.
- [100] Matthew Greenberg. Computing Heegner points arising from Shimura curve parametrizations. *Preprint*, 11 pages, 2006.
- [101] Matthew Greenberg. Heegner point computations via numerical p -adic integration. In *Algorithmic Number Theory*, volume 4076 of *Lecture Notes in Computer Science*, pages 361–376. Springer Berlin / Heidelberg, 2006.
- [102] Matthew Greenberg. *Heegner Points and Rigid Analytic Modular Forms*. PhD Thesis, McGill University, 2006.
- [103] Grigor Grigorov, Andrei Jorza, Stephan Patrikis, William A. Stein, and Corina Tarnita-Patrascu. Verification of the Birch and Swinnerton-Dyer conjecture for specific elliptic curves. *Preprint*, 26 pages.
- [104] Jordi Guàrdia. Jacobian Nullwerte, periods and symmetric equations for hyperelliptic curves. *Ann. Inst. Fourier (Grenoble)*, 57(4):1253–1283, 2007.
- [105] Brian Hansen. *Explicit computations supporting a generalization of Serre’s conjecture*. Master of Science thesis, Brigham Young University, 2005.

- [106] Robin Hartshorne and Ronald van Luijk. Non-Euclidean Pythagorean triples, a problem of Euler, and rational points on $K3$ surfaces. *Math. Intelligencer*, 30(4):4–10, 2008.
- [107] Ki-ichiro Hashimoto, Katsuya Miyake, and Hiroaki Nakamura, editors. *Galois Theory and Modular Forms*, volume 11 of *Developments in Mathematics*, Boston, MA, 2004. Kluwer Academic Publishers.
- [108] Florian Hess. Computing relations in divisor class groups of algebraic curves over finite fields. *Preprint*, 2003.
- [109] Florian Hess. A note on the Tate pairing of curves over finite fields. *Arch. Math. (Basel)*, 82(1):28–32, 2004.
- [110] Laura Hitt. Families of genus 2 curves with small embedding degree. *J. Math. Cryptol.*, 3(1):19–36, 2009.
- [111] E. W. Howe and K. E. Lauter. Improved upper bounds for the number of points on curves over finite fields. *Ann. Inst. Fourier (Grenoble)*, 53(6):1677–1737, 2003.
- [112] Everett W. Howe. Infinite families of pairs of curves over Q with isomorphic Jacobians. *J. London Math. Soc. (2)*, 72(2):327–350, 2005.
- [113] Everett W. Howe. Supersingular genus-2 curves over fields of characteristic 3. In *Computational arithmetic geometry*, volume 463 of *Contemp. Math.*, pages 49–69. Amer. Math. Soc., Providence, RI, 2008.
- [114] Everett W. Howe, Kristin E. Lauter, and Jaap Top. Pointless curves of genus three and four. In *Arithmetic, Geometry and Coding Theory (AGCT 2003)*, volume 11 of *Sémin. Congr.*, pages 125–141. Soc. Math. France, Paris, 2005.
- [115] Everett W. Howe and Hui June Zhu. On the existence of absolutely simple abelian varieties of a given dimension over an arbitrary field. *J. Number Theory*, 92(1):139–163, 2002.
- [116] Hendrik Hubrechts. Point counting in families of hyperelliptic curves. *Found. Comput. Math.*, 8(1):137–169, 2008.
- [117] Hendrik Hubrechts. Quasi-quadratic elliptic curve point counting using rigid cohomology. *J. Symb. Comput.*, 44(9):1255–1267, 2009.

- [118] Klaus Hulek and Helena Verrill. On modularity of rigid and nonrigid Calabi-Yau varieties associated to the root lattice A_4 . *Nagoya Math. J.*, 179:103–146, 2005.
- [119] Klaus Hulek and Helena A. Verrill. On the motive of Kummer varieties associated to $\Gamma_1(7)$ — Supplement to the paper: “The modularity of certain non-rigid Calabi-Yau threefolds” by R. Livné and N. Yui. *J. Math. Kyoto Univ.*, 45(4):667–681, 2005.
- [120] Patrick Ingram. Multiples of integral points on elliptic curves. *Journal of Number Theory*, 129(1):182 – 208, 2009.
- [121] Farzali A. Izadi and V. Kumar Murty. Counting points on an abelian variety over a finite field. In *Progress in Cryptology—Indocrypt 2003*, volume 2904 of *Lecture Notes in Comput. Sci.*, pages 323–333. Springer, Berlin, 2003.
- [122] Dimitar Jetchev, Kristin Lauter, and William Stein. Explicit Heegner points: Kolyvagin’s conjecture and non-trivial elements in the Shafarevich-Tate group. *Journal of Number Theory*, 129(2):284 – 302, 2009.
- [123] Dimitar P. Jetchev and William A. Stein. Visibility of the Shafarevich-Tate group at higher level. *Doc. Math.*, 12:673–696, 2007.
- [124] Jorge Jimenez-Urroz and Tonghai Yang. Heegner zeros of theta functions. *Trans. Amer. Math. Soc.*, 355(10):4137–4149 (electronic), 2003.
- [125] David Joyner and Amy Ksir. Modular representations on some Riemann-Roch spaces of modular curves $X(N)$. In *Computational Aspects of Algebraic Curves*, volume 13 of *Lecture Notes Ser. Comput.*, pages 163–205. World Sci. Publ., Hackensack, NJ, 2005.
- [126] Koray Karabina and Edlyn Teske. On prime-order elliptic curves with embedding degrees $k=3,4$, and 6. In *Algorithmic Number Theory*, volume 5011 of *Lecture Notes in Computer Science*, pages 102–117. Springer, 2008.
- [127] L. J. P. Kilford. Some non-Gorenstein Hecke algebras attached to spaces of modular forms. *J. Number Theory*, 97(1):157–164, 2002.

- [128] L. J. P. Kilford. Slopes of 2-adic overconvergent modular forms with small level. *Math. Res. Lett.*, 11(5-6):723–739, 2004.
- [129] L. J. P. Kilford. On a p -adic extension of the Jacquet-Langlands correspondence to weight 1. [arXiv:0809.1048v1](https://arxiv.org/abs/0809.1048v1) [math.NT], 17 pages, 2008.
- [130] David R. Kohel. Hecke module structure of quaternions. In *Class Field Theory—Its Centenary and Prospect (Tokyo, 1998)*, volume 30 of *Adv. Stud. Pure Math.*, pages 177–195. Math. Soc. Japan, Tokyo, 2001.
- [131] David R. Kohel. The AGM- $X_0(N)$ Heegner point lifting algorithm and elliptic curve point counting. In *Advances in Cryptology—Asiacrypt 2003*, volume 2894 of *Lecture Notes in Comput. Sci.*, pages 124–136. Springer, Berlin, 2003.
- [132] David R. Kohel and William A. Stein. Component groups of quotients of $J_0(N)$. In *Algorithmic Number Theory (Leiden, 2000)*, volume 1838 of *Lecture Notes in Comput. Sci.*, pages 405–412. Springer, Berlin, 2000.
- [133] David R. Kohel and Helena A. Verrill. Fundamental domains for Shimura curves. *J. Théor. Nombres Bordeaux*, 15(1):205–222, 2003.
- [134] Kenji Koike and Annegret Weng. Construction of CM Picard curves. *Math. Comp.*, 74(249):499–518 (electronic), 2005.
- [135] Elisavet Konstantinou and Kontogeorgis Aristides. Computing polynomials of the Ramanujan \mathfrak{t}_n class invariants. *Canad. Math. Bull.*, To appear, 12 pages, 2007.
- [136] Aristides Kontogeorgis and Victor Rotger. On the non-existence of exceptional automorphisms on Shimura curves. *Bull. Lond. Math. Soc.*, 40(3):363–374, 2008.
- [137] Andrew Kresch and Yuri Tschinkel. Integral points on punctured abelian surfaces. In *Algorithmic Number Theory (Sydney, 2002)*, volume 2369 of *Lecture Notes in Comput. Sci.*, pages 198–204. Springer, Berlin, 2002.

- [138] L. Kulesz, G. Matera, and É. Schost. Uniform bounds on the number of rational points of a family of curves of genus 2. *J. Number Theory*, 108(2):241–267, 2004.
- [139] Dominic Lanphier. The trace of special values of modular L -functions. *Preprint*, 18 pages.
- [140] Alan G. B. Lauder. Ranks of elliptic curves over function fields. *LMS J. Comput. Math.*, 11:172–212, 2008.
- [141] F. Leprévost, M. Pohst, and A. Schöpp. Rational torsion of $J_0(N)$ for hyperelliptic modular curves and families of Jacobians of genus 2 and genus 3 curves with a rational point of order 5, 7 or 10. *Abh. Math. Sem. Univ. Hamburg*, 74:193–203, 2004.
- [142] Franck Leprévost, Michael Pohst, and Andreas Schöpp. Familles de polynômes liées aux courbes modulaires $X(l)$ unicursales et points rationnels non-triviaux de courbes elliptiques quotient. *Acta Arith.*, 110(4):401–410, 2003.
- [143] Reynald Lercier and David Lubicz. A quasi-quadratic time algorithm for hyperelliptic curve point counting. *The Ramanujan Journal*, 12(3):399–423, 2006.
- [144] Reynald Lercier and Thomas Sirvent. On Elkies subgroups of l -torsion points in elliptic curves defined over a finite field. *J. Théor. Nombres Bordeaux*, 20(3):783–797, 2008.
- [145] David Loeffler. Explicit calculations of automorphic forms for definite unitary groups. *LMS J. Comput. Math.*, 11:326–342, 2008.
- [146] Adam Logan and Ronald van Luijk. Nontrivial elements of Sha explained through $K3$ surfaces. *Math. Comp.*, 78(265):441–483, 2009.
- [147] Ling Long and Chris Kurth. On modular forms for some noncongruence subgroups of $SL_2\mathbb{Z}$ II. [arXiv:0809.0020v1](https://arxiv.org/abs/0809.0020v1) [[math.NT](https://arxiv.org/abs/0809.0020v1)], 13 pages, 2008.
- [148] Dino Lorenzini and Thomas J. Tucker. Thue equations and the method of Coleman-Chabauty. [arXiv:math.NT/0005186](https://arxiv.org/abs/math.NT/0005186), 30 pages, 2000.

- [149] Kazuo Matsuno. Construction of elliptic curves with large Iwasawa λ -invariants and large Tate-Shafarevich groups. *Manuscripta Math.*, 122(3):289–304, 2007.
- [150] Kazuto Matsuo, Jinhui Chao, and Shigeo Tsujii. An improved baby step giant step algorithm for point counting of hyperelliptic curves over finite fields. In *Algorithmic Number Theory (Sydney, 2002)*, volume 2369 of *Lecture Notes in Comput. Sci.*, pages 461–474. Springer, Berlin, 2002.
- [151] J. Miret, R. Moreno, A. Rio, and M. Valls. Computing the l -power torsion of an elliptic curve over a finite field. *Math. Comp.*, 78(267):1767–1786, 2009.
- [152] J. Miret, R. Moreno, D. Sadornil, J. Tena, and M. Valls. Computing the height of volcanoes of l -isogenies of elliptic curves over finite fields. *Appl. Math. Comput.*, 196(1):67–76, 2008.
- [153] Annika Niehage. *Quantum Goppa Codes over Hyperelliptic Curves*. Diplomarbeit, Universität Mannheim, 2004.
- [154] Mihran Papikian. On the variation of Tate-Shafarevich groups of elliptic curves over hyperelliptic curves. *J. Number Theory*, 115(2):249–283, 2005.
- [155] Bernadette Perrin-Riou. Arithmétique des courbes elliptiques à réduction supersingulière en p . *Experiment. Math.*, 12(2):155–186, 2003.
- [156] Bjorn Poonen. Computational aspects of curves of genus at least 2. In *Algorithmic Number Theory (Talence, 1996)*, volume 1122 of *Lecture Notes in Comput. Sci.*, pages 283–306. Springer, Berlin, 1996.
- [157] Bjorn Poonen, Edward F. Schaefer, and Michael Stoll. Twists of $X(7)$ and primitive solutions to $x^2 + y^3 = z^7$. *Duke Math. J.*, 137(1):103–158, 2007.
- [158] Lisa Marie Redekop. *Torsion Points of Low Order on Elliptic Curves and Drinfeld Modules*. PhD thesis, 2002.

- [159] Guillaume Ricotta and Thomas Vidick. Hauteur asymptotique des points de Heegner. *Canad. J. Math.*, 60(6):1406–1436, 2008.
- [160] Christophe Ritzenthaler. Automorphismes des courbes modulaires $X(n)$ en caractéristique p . *Manuscripta Math.*, 109(1):49–62, 2002.
- [161] Christophe Ritzenthaler. Point counting on genus 3 non hyperelliptic curves. In *Algorithmic Number Theory*, volume 3076 of *Lecture Notes in Comput. Sci.*, pages 379–394. Springer, Berlin, 2004.
- [162] David Savitt. The maximum number of points on a curve of genus 4 over F_8 is 25. *Canad. J. Math.*, 55(2):331–352, 2003.
- [163] Edward F. Schaefer and Michael Stoll. How to do a p -descent on an elliptic curve. *Trans. Amer. Math. Soc.*, 356(3):1209–1231 (electronic), 2004.
- [164] Jasper Scholten. Weil restriction of an elliptic curve over a quadratic extension. *Preprint*, 6 pages, 2004.
- [165] Andreas M. Schöpp. *Über Torsionspunkte elliptischer und hyperelliptischer Kurven nebst Anwendungen*. PhD thesis, Technische Universität Berlin,, April 2005.
- [166] Samir Siksek. On standardized models of isogenous elliptic curves. *Math. Comp.*, 74(250):949–951 (electronic), 2005.
- [167] Samir Siksek and John E. Cremona. On the Diophantine equation $x^2 + 7 = y^m$. *Acta Arith.*, 109(2):143–149, 2003.
- [168] Benjamin Smith. Isogenies and the discrete logarithm problem in Jacobians of genus 3 hyperelliptic curves. In *Advances in Cryptology, Eurocrypt 2008*, volume 4965 of *Lecture Notes in Computer Science*, pages 163–180. Springer Berlin/Heidelberg, 2008.
- [169] Sebastian Karl Michael Stamminger. *Explicit 8-Descent on Elliptic Curves*. PhD thesis, International University Bremen, 2005.
- [170] William Stein. Studying the Birch and Swinnerton-Dyer conjecture for modular abelian varieties using Magma. In *Discovering Mathematics with Magma*, volume 19 of *Algorithms Comput. Math.*, pages 93–116. Springer, Berlin, 2006.

- [171] William A. Stein. There are genus one curves over Q of every odd index. *J. Reine Angew. Math.*, 547:139–147, 2002.
- [172] William A. Stein. Shafarevich-Tate groups of nonsquare order. In *Modular curves and abelian varieties*, volume 224 of *Progr. Math.*, pages 277–289. Birkhäuser, Basel, 2004.
- [173] William A. Stein. Visibility of Mordell-Weil groups. *Doc. Math.*, 12:587–606, 2007.
- [174] Michael Stoll. Implementing 2-descent for Jacobians of hyperelliptic curves. *Acta Arith.*, 98(3):245–277, 2001.
- [175] Michael Stoll. On the height constant for curves of genus two. II. *Acta Arith.*, 104(2):165–182, 2002.
- [176] Michael Stoll. Rational 6-cycles under iteration of quadratic polynomials. *LMS J. Comput. Math.*, 11:367–380, 2008.
- [177] Thotsaphon Thongjunthug. Computing a lower bound for the canonical height on elliptic curves over totally real number fields. In *Algorithmic Number Theory*, volume 5011 of *Lecture Notes in Computer Science*, pages 139–152. Springer, 2008.
- [178] Hans-Christian Graf v. Bothmer. Finite field experiments (with an appendix by Stefan Wiedmann). In *Higher-Dimensional Geometry over Finite Fields*, volume 16 of *NATO Science for Peace and Security Series, D: Information and Communication Security*, pages 1–62. 2008.
- [179] Marie-France Vignéras. p -adic integral structures of some representations of $GL(2, F)$. *Preprint*, 23 pages, 2005.
- [180] Bogdan G. Vioreanu. Mordell-Weil problem for cubic surfaces, numerical evidence. [arXiv:0802.0742v1](https://arxiv.org/abs/0802.0742v1) [[math.AG](https://arxiv.org/abs/0802.0742v1)], 22 pages, 2008.
- [181] Mark Watkins. A note on integral points on elliptic curves. *J. Théor. Nombres Bordeaux*, 18(3):707–720, 2006.
- [182] Mark Watkins. Some remarks on Heegner point computations. [arXiv:math.NT/0506325](https://arxiv.org/abs/math.NT/0506325), 14 pages, 2006.

- [183] Mark Watkins. Some heuristics about elliptic curves. *Experiment. Math.*, 17(1):105–125, 2008.
- [184] Rolf Stefan Wilke. On rational embeddings of curves in the second Garcia-Stichtenoth tower. *Finite Fields Appl.*, 14(2):494–504, 2008.
- [185] Christian Wuthrich. The fine Tate-Shafarevich group. *Math. Proc. Cambridge Philos. Soc.*, 142(1):1–12, 2007.
- [186] Chaoping Xing. Applications of algebraic curves to constructions of sequences. In *Cryptography and Computational Number Theory (Singapore, 1999)*, volume 20 of *Progr. Comput. Sci. Appl. Logic*, pages 137–146. Birkhäuser, Basel, 2001.
- [187] Chaoping Xing and Sze Ling Yeo. Construction of global function fields from linear codes and vice versa. *Trans. Amer. Math. Soc.*, 361(3):1333–1349, 2008.
- [188] Huilin Zhu and Jianhua Chen. Integral points on a class of elliptic curve. *Wuhan Univ. J. Nat. Sci.*, 11(3):477–480, 2006.