

Combinatorics

Designs and Configurations

05Bxx

- [1] David Applegate, E. M. Rains, and N. J. A. Sloane. On asymmetric coverings and covering numbers. *J. Combin. Des.*, 11(3):218–228, 2003.
- [2] Makoto Araya, Masaaki Harada, and Hadi Kharaghani. Some Hadamard matrices of order 32 and their binary codes. *J. Combin. Des.*, 12(2):142–146, 2004.
- [3] R. A. Bailey, Peter J. Cameron, Peter Dobcsányi, John P. Morgan, and Leonard H. Soicher. Designs on the web. *Discrete Math.*, 306(23):3014–3027, 2006.
- [4] John Bamberg and Tim Penttila. A classification of transitive ovoids, spreads, and m -systems of polar spaces. *Forum Math.*, 21(2):181–216, 2009.
- [5] Eiichi Bannai and Etsuko Bannai. On Euclidean tight 4-designs. *J. Math. Soc. Japan*, 58(3):775–804, 2006.
- [6] L. M. Batten and J. M. Dover. Some sets of type (m, n) in cubic order planes. *Des. Codes Cryptogr.*, 16(3):211–213, 1999.
- [7] Thomas Beth, Christopher Charney, Markus Grassl, Gernot Alber, Aldo Delgado, and Michael Mussinger. A new class of designs which protect against quantum jumps. *Des. Codes Cryptogr.*, 29(1-3):51–70, 2003.
- [8] Anton Betten, Adalbert Kerber, Reinhard Laue, and Alfred Wassermann. Simple 8-designs with small parameters. *Des. Codes Cryptogr.*, 15(1):5–27, 1998.
- [9] A. Bonnet, E. Rains, and P. Solé. 3-colored 5-designs and Z_4 -codes. *J. Statist. Plann. Inference*, 86(2):349–368, 2000.
- [10] A. Bonnet, P. Solé, and P. Udaya. Tricolore 3-designs in type III codes. *Discrete Math.*, 241(1-3):129–138, 2001.

- [11] Alexis Bonnetcaze, Bernard Mourrain, and Patrick Solé. Jacobi polynomials, type II codes, and designs. *Des. Codes Cryptogr.*, 16(3):215–234, 1999.
- [12] Thomas Britz and Carrie G. Rutherford. Covering radii are not matroid invariants. *Discrete Math.*, 296(1):117–120, 2005.
- [13] Thomas Britz and Keisuke Shiromoto. Designs from subcode supports of linear codes. *Des. Codes Cryptogr.*, 46(2):175–189, 2008.
- [14] S. Allen Broughton, Robert M. Dirks, Maria T. Slougher, and C. Ryan Vinroot. Triangular surface tiling groups for low genus. *Preprint*, 2001.
- [15] I. Cardinali, N. Durante, T. Penttila, and R. Trombetti. Bruen chains over fields of small order. *Discrete Math.*, 282(1-3):245–247, 2004.
- [16] L. L. Carpenter and J. D. Key. On Hadamard matrices from resolvable Steiner designs. In *Proceedings of the Twenty-sixth Southeastern International Conference on Combinatorics, Graph Theory and Computing (Boca Raton, FL, 1995)*, volume 108, pages 53–63, 1995.
- [17] David B. Chandler and Qing Xiang. Cyclic relative difference sets and their p -ranks. *Des. Codes Cryptogr.*, 30(3):325–343, 2003.
- [18] Chris Charnes, Martin Rötteler, and Thomas Beth. Homogeneous bent functions, invariants, and designs. *Des. Codes Cryptogr.*, 26(1-3):139–154, 2002.
- [19] D. Combe, W. D. Palmer, and W. R. Unger. Bhaskar Rao designs and the alternating group A_4 . *Australas. J. Combin.*, 24:275–283, 2001.
- [20] Marston Conder and John McKay. Markings of the Golay code. *New Zealand J. Math.*, 25(2):133–139, 1996.
- [21] Antonio Cossidente and Sam K. J. Vereecke. Some geometry of the isomorphism $\text{Sp}(4, q) \cong \text{O}(5, q)$, q even. *J. Geom.*, 70(1-2):28–37, 2001.
- [22] Warwick de Launey and Richard M. Stafford. On cocyclic weighing matrices and the regular group actions of certain Paley matrices. *Discrete Appl. Math.*, 102(1-2):63–101, 2000.

- [23] U. Dempwolff. Automorphisms and equivalence of bent functions and of difference sets in elementary abelian 2-groups. *Comm. Algebra*, 34(3):1077–1131, 2006.
- [24] Cunsheng Ding, Zeying Wang, and Qing Xiang. Skew Hadamard difference sets from the Ree-Tits slice symplectic spreads in $\text{PG}(3, 3^{2h+1})$. *J. Combin. Theory Ser. A*, 114(5):867–887, 2007.
- [25] Cunsheng Ding and Jin Yuan. A family of skew Hadamard difference sets. *J. Combin. Theory Ser. A*, 113(7):1526–1535, 2006.
- [26] G. L. Ebert. Quasimultiples of geometric designs. *Discrete Math.*, 284(1-3):123–129, 2004.
- [27] Ronald Evans, Henk D. L. Hollmann, Christian Krattenthaler, and Qing Xiang. Gauss sums, Jacobi sums, and p -ranks of cyclic difference sets. *J. Combin. Theory Ser. A*, 87(1):74–119, 1999.
- [28] W. Fish, J. D. Key, and E. Mwambene. Graphs, designs and codes related to the n -cube. *Discrete Math.*, 309(10):3255–3269, 2009.
- [29] D. L. Flannery. Cocyclic Hadamard matrices and Hadamard groups are equivalent. *J. Algebra*, 192(2):749–779, 1997.
- [30] Anna Fukshansky and Corinna Wiedorn. C -extensions of the Petersen geometry for M_{22} . *European J. Combin.*, 20(3):233–238, 1999.
- [31] S. Georgiou, I. Kotsireas, and C. Koukouvinos. Inequivalent Hadamard matrices of order $2n$ constructed from Hadamard matrices of order n . *J. Combin. Math. Combin. Comput.*, 63:65–79, 2007.
- [32] S. Georgiou and C. Koukouvinos. Some results on orthogonal designs and Hadamard matrices. *Int. J. Appl. Math.*, 17(4):433–443, 2005.
- [33] Stelios D. Georgiou. New two-variable full orthogonal designs and related experiments with linear regression models. *Statist. Probab. Lett.*, 77(1):25–31, 2007.
- [34] Claudia Gohlich, Helmut Koch, and Gabriele Nebe. Block squares. *Math. Nachr.*, 241:73–102, 2002.

- [35] Masaaki Harada. On the self-dual F_5 -codes constructed from Hadamard matrices of order 24. *J. Combin. Des.*, 13(2):152–156, 2005.
- [36] Masaaki Harada. Self-orthogonal 3-(56, 12, 65) designs and extremal doubly-even self-dual codes of length 56. *Des. Codes Cryptogr.*, 38(1):5–16, 2006.
- [37] Masaaki Harada, Clement Lam, and Vladimir D. Tonchev. Symmetric (4, 4)-nets and generalized Hadamard matrices over groups of order 4. *Des. Codes Cryptogr.*, 34(1):71–87, 2005.
- [38] Masaaki Harada and Akihiro Munemasa. A quasi-symmetric 2-(49, 9, 6) design. *J. Combin. Des.*, 10(3):173–179, 2002.
- [39] Masaaki Harada and Vladimir D. Tonchev. Self-orthogonal codes from symmetric designs with fixed-point-free automorphisms. *Discrete Math.*, 264(1-3):81–90, 2003.
- [40] K. J. Horadam. An introduction to cocyclic generalised Hadamard matrices. *Discrete Appl. Math.*, 102(1-2):115–131, 2000.
- [41] Naoyuki Horiguchi, Hiroyuki Nakasora, and Takehisa Wakabayashi. On the strongly regular graphs obtained from quasi-symmetric 2-(31, 7, 7) designs. *Bull. Yamagata Univ. Natur. Sci.*, 16(1):1–6, 2005.
- [42] Masakazu Jimbo and Keisuke Shiromoto. A construction of mutually disjoint Steiner systems from isomorphic Golay codes. *J. Combin. Theory Ser. A*, 116(7):1245–1251, 2009.
- [43] J. D. Key and M. J. de Resmini. Small sets of even type and codewords. *J. Geom.*, 61(1-2):83–104, 1998.
- [44] J. D. Key and K. Mackenzie-Fleming. Rigidity theorems for a class of affine resolvable designs. *J. Combin. Math. Combin. Comput.*, 35:147–160, 2000.
- [45] J. D. Key, T. P. McDonough, and V. C. Mavron. Partial permutation decoding for codes from finite planes. *European J. Combin.*, 26(5):665–682, 2005.

- [46] J. D. Key and J. Moori. Codes, designs and graphs from the Janko groups J_1 and J_2 . *J. Combin. Math. Combin. Comput.*, 40:143–159, 2002.
- [47] J. D. Key and F. D. Shobe. Some transitive Steiner triple systems of Bagchi and Bagchi. *J. Statist. Plann. Inference*, 58(1):79–86, 1997.
- [48] J. D. Key and F. E. Sullivan. Steiner triple systems with many affine hyperplanes. In *Proceedings of the Twenty-sixth Southeastern International Conference on Combinatorics, Graph Theory and Computing (Boca Raton, FL, 1995)*, volume 107, pages 105–112, 1995.
- [49] Michael Kiermaier and Sascha Kurz. Maximal integral point sets in affine planes over finite fields. *Discrete Math.*, 309(13):4564–4575, 2009.
- [50] Jon-Lark Kim and Vera Pless. Designs in additive codes over $\text{GF}(4)$. *Des. Codes Cryptogr.*, 30(2):187–199, 2003.
- [51] Jon-Lark Kim and Patrick Solé. Skew Hadamard designs and their codes. *Des. Codes Cryptogr.*, 49(1-3):135–145, 2008.
- [52] I. S. Kotsireas and C. Koukouvinos. Inequivalent hadamard matrices from orthogonal designs. In *PASCO '07: Proceedings of the 2007 International Workshop on Parallel Symbolic Computation*, pages 95–96, New York, NY, USA, 2007. ACM.
- [53] Ilias S. Kotsireas and Christos Koukouvinos. Constructions for Hadamard matrices of Williamson type. *J. Combin. Math. Combin. Comput.*, 59:17–32, 2006.
- [54] Ilias S. Kotsireas and Christos Koukouvinos. Orthogonal designs via computational algebra. *J. Combin. Des.*, 14(5):351–362, 2006.
- [55] Ilias S. Kotsireas, Christos Koukouvinos, and Jennifer Seberry. Hadamard ideals and Hadamard matrices with circulant core. *J. Combin. Math. Combin. Comput.*, 57:47–63, 2006.
- [56] Ilias S. Kotsireas, Christos Koukouvinos, and Jennifer Seberry. Hadamard ideals and Hadamard matrices with two circulant cores. *European J. Combin.*, 27(5):658–668, 2006.

- [57] C. Koukouvinos and S. Stylianou. On skew-Hadamard matrices. *Discrete Math.*, 308(13):2723–2731, 2008.
- [58] Christos Koukouvinos and Dimitris E. Simos. Improving the lower bounds on inequivalent Hadamard matrices through orthogonal designs and meta-programming techniques. *Applied Numerical Mathematics*, In Press, 2009.
- [59] Wolfgang Lempken. Two new symmetric 2-(144, 66, 30) designs. *Preprint*.
- [60] Keith E. Mellinger. Designs, Geometry, and a Golfer’s Dilemma. *Math. Mag.*, 77(4):275–282, 2004.
- [61] Jamshid Moori and B. G. Rodrigues. Some designs and codes invariant under the simple group Co_2 . *J. Algebra*, 316(2):649–661, 2007.
- [62] Akihiro Munemasa and Vladimir D. Tonchev. A new quasi-symmetric 2-(56, 16, 6) design obtained from codes. *Discrete Math.*, 284(1-3):231–234, 2004.
- [63] Christopher Parker and Vladimir D. Tonchev. Linear codes and doubly transitive symmetric designs. *Linear Algebra Appl.*, 226/228:237–246, 1995.
- [64] E. M. Rains, N. J. A. Sloane, and John Stufken. The lattice of N -run orthogonal arrays. *J. Statist. Plann. Inference*, 102(2):477–500, 2002.
- [65] Colin Ramsay. *Trades and Defining Sets: Theoretical and Computational Results*. PhD thesis, The University of Queensland, 1998.
- [66] B. G. Rodrigues. Self-orthogonal designs and codes from the symplectic groups $S_4(3)$ and $S_4(4)$. *Discrete Math.*, 308(10):1941–1950, 2008.
- [67] Chekad Sarami and Vladimir D. Tonchev. Cyclic quasi-symmetric designs and self-orthogonal codes of length 63. *J. Statist. Plann. Inference*, 138(1):80–85, 2008.
- [68] Michel Sebillé. On a result of Cameron and Praeger on block-transitive point-imprimitive t -designs. In *Algebraic Combinatorics and Applications (Gößweinstein, 1999)*, pages 316–323. Springer, Berlin, 2001.

- [69] Junichi Shigezumi. On 3-lattices and spherical designs. arXiv:0810.4373, 199 pages, 2008.
- [70] Pawel Wocjan. Efficient decoupling schemes with bounded controls based on Eulerian orthogonal arrays. *Phy. Rev. A.*, 73(6):7, 2006.