Computational Methods

12-04

[1] Gregory V. Bard, Nicolas T. Courtois, and Chris Jefferson, *Efficient methods for conversion and solution of sparse systems of low-degree multivariate polynomials over GF(2) via SAT-solvers*, 2007.

[2] Cristina Bertone, Guillaume Chéze, and André Galligo, *Modular Las Vegas algorithms for polynomial absolute factorization*, J. Symbolic Comput. **45** (2010), no. 12, 1280–1295.

[3] Thomas Beth, Jörn Müller-Quade, and Rainer Steinwandt, *Computing restrictions of ideals in finitely generated k-algebras by means of Buchberger's algorithm*, J. Symbolic Comput. **41** (2006), no. 3-4, 372–380. MR MR2202557 (2006j:13027)

[4] A. Bostan, G. Lecerf, B. Salvy, É. Schost, and B. Wiebelt, *Complexity issues in bivariate polynomial factorization*, ISSAC '04: Proceedings of the 2004 International Symposium on Symbolic and Algebraic Computation (New York, NY, USA), ACM Press, 2004, pp. 42–49.

[5] Guillaume Chèze and Grégoire Lecerf, *Lifting and recombination techniques for absolute factorization*, J. Complexity **23** (2007), no. 3, 380–420. MR MR2330992

[6] Akpodigha Filatei, Xin Li, Marc Moreno Maza, and Éric Schost, *Implementation techniques for fast polynomial arithmetic in a high-level programming environment*, ISSAC '06: Proceedings of the 2006 International Symposium on Symbolic and Algebraic Computation (New York, NY, USA), ACM Press, 2006, pp. 93–100.

[7] Katharina Geißler and Jürgen Klüners, *Galois group computation for rational polynomials*, J. Symbolic Comput. **30** (2000), no. 6, 653–674, Algorithmic methods in Galois theory. MR MR1800032 (2001k:12006)

[8] Kiran S. Kedlaya, *Search techniques for root-unitary polynomials*, Computational arithmetic geometry, Contemp. Math., vol. 463, Amer. Math. Soc., Providence, RI, 2008, pp. 71–81. MR MR2459990 (2009h:26022)

[9] Sara Khodadad and Michael Monagan, *Fast rational function reconstruction*, ISSAC 2006, ACM, New York, 2006, pp. 184–190. MR MR2289118

[10] Grégoire Lecerf, *New recombination algorithms for bivariate polynomial factorization based on Hensel lifting*, Appl. Algebra Engrg. Comm. Comput. **21** (2010), no. 2, 151–176. MR 2600710

[11] Hsin-Chao Liao and Richard J. Fateman, *Evaluation of the heuristic polynomial GCD*, ISSAC '95: Proceedings of the 1995 International Symposium on Symbolic and Algebraic Computation (New York, NY, USA), ACM Press, 1995, pp. 240–247.

[12] Michael Monagan, *Maximal quotient rational reconstruction: An almost optimal algorithm for rational reconstruction*, ISSAC 2004, ACM, New York, 2004, pp. 243–249. MR MR2126950 (2005j:68137)

[13] Jörn Müller-Quade and Rainer Steinwandt, *Basic algorithms for rational function fields*, J. Symbolic Comput. **27** (1999), no. 2, 143–170. MR MR1672124 (2000a:13043)

[14] ———, *Gröbner bases applied to finitely generated field extensions*, J. Symbolic Comput. **30** (2000), no. 4, 469–490. MR MR1784753 (2001i:13040)

[15] Fatima K. Abu Salem and Rawan N. Soudah, *An empirical study of cache-oblivious polygon indecomposability testing*, Computing **88** (2010), no. 8, 55–78.

[16] Leonard Soicher and John McKay, *Computing Galois groups over the rationals*, J. Number Theory **20** (1985), no. 3, 273–281. MR MR797178 (87a:12002)

[17] Allan Steel, *A new scheme for computing with algebraically closed fields*, Algorithmic Number Theory (Sydney, 2002), Lecture Notes in Comput. Sci., vol. 2369, Springer, Berlin, 2002, pp. 491–505. MR MR2041106 (2005b:12016)

[18] Allan K. Steel, *Computing with algebraically closed fields*, J. Symbolic Comput. **45** (2010), no. 3, 342–372.

[19] Rainer Steinwandt, *On computing a separating transcendence basis*, SIGSAM Bulletin **34** (2000), no. 4.